

Some recent results on evaluating Feynman integrals

V.A. Smirnov^{a*}

^a Nuclear Physics Institute of Moscow State University,
Moscow 119992, Russia

Some recent results on evaluating Feynman integrals are reviewed. The status of the method based on Mellin-Barnes representation as a powerful tool to evaluate individual Feynman integrals is characterized. A new method based on Gröbner bases to solve integration by parts relations in an automatic way is described.

1. Introduction

Perturbative quantum-theoretical amplitudes are expressed in terms of Feynman integrals over loop momenta. Usually, one turns immediately to scalar Feynman integrals using some projectors and obtains a family of scalar Feynman integrals with the same structure of the integrand and various powers of propagators (indices):

$$F(a_1, \dots, a_n) = \int \dots \int \frac{d^d k_1 \dots d^d k_h}{E_1^{a_1} \dots E_n^{a_n}}, \quad (1)$$

where k_i , $i = 1, \dots, h$, are loop momenta and the denominators E_r are either quadratic or linear with respect to k_i and external momenta $q_1, \dots, q_N$. By default, the integrals are dimensionally regularized with $d = 4 - 2\epsilon$.

If the number of Feynman integrals needed for a given calculation is small or/and they are simple, one evaluates, by some methods, every scalar Feynman integral of the given family. Various methods are used, in particular, alpha and/or Feynman parameters, Mellin-Barnes (MB) representation [1,2] and differential equations [3]. In the next section, the method of MB representation is briefly reviewed.

If it is necessary to evaluate a lot of complicated Feynman integrals (1) the standard way is to apply integration by parts (IBP) [4] relations

$$\int d^d k_1 \int d^d k_2 \dots \frac{\partial}{\partial k_i} \left(p_j \frac{1}{E_1^{a_1} \dots E_n^{a_n}} \right) = 0$$

*Supported by the Russian Foundation for Basic Research through grant 05-02-17645. To appear in the proceedings of the 7th International Symposium on Radiative Corrections (RADCOR05), Shonan Village, Japan, 2005.

in order to construct an algorithm that gives and expression of a given Feynman integral as a linear combination of some *master* integrals, $F(a_1, \dots, a_n) = \sum_i c(a_1, \dots, a_n) I_i$. Here $p_1 = k_1, \dots, p_h = k_h, p_{h+1} = q_1, \dots, p_{h+N} = q_N$. Once one has a solution of this (reduction) problem, it becomes sufficient to evaluate only master integrals. The IBP relations can be written as $f_i \cdot F$ where f_i are polynomials in the shift operators $Y_i^{\pm 1}$ defined by $(Y_i^{\pm 1} \cdot F)(a_1, \dots, a_n) = F(a_1, \dots, a_i \pm 1, \dots, a_n)$.

The first attempt to make the reduction procedure systematic was based on the fact that the total number of IBP equations grows faster than the number of Feynman integrals satisfying the condition $\sum_i |a_i| \leq M$ at $M \rightarrow \infty$ so that, at sufficiently large M , one obtains an overdetermined system of equations [5,6] which can be solved. There is already a public implementation of this algorithm on a computer [7]. Another attempt [8] is based on a special parametric representation.

Other attempts are based on the use of Gröbner bases (GB) [9]. This idea was first suggested in [10], where IBP relations were reduced to differential equations. To do this, it is assumed that there is a non-zero mass for each line. The typical combination $a_i Y_i$, is then naturally transformed into the operator of differentiation in the corresponding mass. An attempt to use GB associated with the shift operators was made in [11]. However, the corresponding algorithms can now work only in simplest cases, for $n = 2$.

In Section 3, another approach [12] based on GB is briefly characterized. In conclusion, I shall discuss some perspectives.


2. Evaluating by MB representation

The method of MB representation was successfully applied to evaluate massless on-shell double [1,2,13,14,15] and triple [16,17] boxes, with results written in terms of harmonic polylogarithms [18], double boxes with one leg off shell [6,19] and massive on-shell double boxes [20,21]. The method is based on the MB representation

$$\frac{1}{(X+Y)^\lambda} = \int_{-i\infty}^{+i\infty} \frac{Y^z}{X^{\lambda+z}} \frac{\Gamma(\lambda+z)\Gamma(-z)}{\Gamma(\lambda)} \frac{dz}{2\pi i}$$

applied to replace a sum of two terms raised to some power by their products in some powers.

Experience shows that a minimal number of MB integrations is achieved if one introduces MB integrations loop by loop, i.e. derives a MB representation for a one-loop subintegral, inserts it into a higher two-loop integral, etc. For example, for the tennis court graph shown in Fig. 1, one can



start from a MB representation for a lower box subintegral with three legs off shell, then insert it into the double-box subintegral to obtain a MB representation for the double box with two (upper) legs off shell, insert it into the whole graph and obtain an eightfold MB representation for the given family of integrals with general powers of the propagators and the power of the numerator $[(l_1 + l_3)^2]^{-a_{11}}$, where $l_{1,3}$ are the momenta flowing through lines 1 and 3 in the same direction:

$$\begin{aligned} & T(a_1, \dots, a_{11}; s, t; \epsilon) \\ &= \frac{(i\pi^{d/2})^3 (-1)^a (-s)^{8-a-3\epsilon}}{\prod_{j=2,4,5,6,7,8} \Gamma(a_j) \Gamma(4 - a_{4567} - 2\epsilon) t^2} \\ &\times \frac{1}{(2\pi i)^8} \int_{-i\infty}^{+i\infty} dw \prod_{j=1}^7 dz_j \left(\frac{t}{s}\right)^w \prod_{j=2}^7 \Gamma(-z_j) \end{aligned}$$

$$\begin{aligned} &\times \frac{\Gamma(a - 8 + 3\epsilon + w) \Gamma(8 - a - 3\epsilon - w)}{\Gamma(a_1 - z_2) \Gamma(a_3 - z_3) \Gamma(a_9 - z_6) \Gamma(a_{10} - z_{47})} \\ &\times \frac{\Gamma(a_5 + z_{14}) \Gamma(a_2 + z_{56}) \Gamma(2 - w + z_5)}{\Gamma(4 - a_{123} - 2\epsilon + z_{123}) \Gamma(8 - a - 4\epsilon - z_5)} \\ &\times \frac{\Gamma(2 - a_{457} - \epsilon - z_{13}) \Gamma(a_{10} - 2 + w - z_{457})}{\Gamma(a_{1234567,11} - 4 + 2\epsilon + z_{4567})} \\ &\times \Gamma(2 - a_{567} - \epsilon - z_{124}) \Gamma(a_{4567} - 2 + \epsilon + z_{1234}) \\ &\times \Gamma(2 - a_{23} - \epsilon + z_{13} - z_5) \Gamma(a_9 - 2 + w - z_{56}) \\ &\times \Gamma(2 - a_{12} - \epsilon + z_{12} - z_{567}) \Gamma(a_7 + z_{123}) \\ &\times \Gamma(a_{123} - 2 + \epsilon - z_{123} + z_{567}) \Gamma(z_{57} - z_1) \\ &\times \Gamma(4 - a_{89,10} - \epsilon - w + z_{4567}), \end{aligned}$$

with $a_{12} = a_1 + a_2$, $a = a_{1\dots,11}$, $z_{14} = z_1 + z_4$, etc.

Deriving MB representations for general indices is very useful because one can apply it for various partial cases and obtain crucial checks. For example, this very representation was used in [17] to calculate $T(1, \dots, 1, -1)$, in a Laurent expansion in ϵ and to check cross order relations in $N = 4$ SUSY gauge theories [22]. Let us stress that one can check such a cumbersome representation in an easy way by considering two partial cases: when one contracts horizontal lines, i.e. in the limit $a_1, a_3, a_4, a_6, a_8 \rightarrow 0$, or vertical lines, i.e. at $a_2, a_5, a_7, a_9, a_{10} \rightarrow 0$. In both cases, one obtains recursively one-loop integrals which can be evaluated in terms of gamma functions for general ϵ . On the other hand, taking such limits reduces to calculating residues in some integration variables. Consider, for example, the limit $a_5 \rightarrow 0$. We have $\Gamma(a_5)$ in the denominator but also the product $\Gamma(a_5 + z_1 + z_4) \Gamma(-z_1) \Gamma(-z_4)$ which is singular in this limit. To reveal the singularity we take residues at $z_1 = 0$ and $z_4 = 0$ and obtain a factor $\Gamma(a_5)$ so that the limit becomes nontrivial.

In the second step, one resolves the singularity structure in ϵ , taking residues and shifting contours, with the goal to obtain a sum of integrals where one can expand integrands in Laurent series in ϵ . One can apply two strategies formulated in [1] and [2]. According to the first strategy, one performs an analysis of the integrand to reveal how poles in ϵ arise. The guiding principle is that the product $\Gamma(a+z)\Gamma(b-z)$, where a and b can depend on the rest of the integration variables, generates, due to the integration over z , the singularity of the type $\Gamma(a+b)$. So, one

thinks of integrations in various orders and then identifies some ‘key’ gamma functions which are crucial for the generation of poles in ϵ . Then one takes residues and shifts contours, starting from first poles of these key gamma functions. For contributions of the residues, the same analysis and procedure is applied. (See [23] for details.)

Within the second strategy [2], one chooses an initial value of ϵ and values of the real parts of the integration variables, $z_i, w, \dots$ in such a way that one can integrate over straight lines. Then one tends ϵ to zero and whenever the real part of the argument of some gamma function vanishes one crosses this pole and adds a corresponding residue which has one integration less and is treated as the initial integral within the same procedure.

The third step of the method is to evaluate integrals expanded in ϵ after the second step. Here one can use Barnes lemmas and their corollaries to perform some of the MB integrations explicitly. In the last integrations which usually carry dependence on the masses and kinematic invariants, one closes contour in the complex plane and sums up corresponding series.

3. Applying Gröbner bases to solve IBP relations

Let $\mathcal{A}$ be the ring of polynomials of n variables $x_1, \dots, x_n$ and $\mathcal{I} \subset \mathcal{A}$ be an ideal with a basis $\{f_1, f_2, \dots, f_k\}^2$. A basis is a *Gröbner basis* if any polynomial $g \in \mathcal{I}$ is *reduced modulo* this basis to zero for any sequence of reductions. To define *reduction* one needs an *ordering of monomials* $cx_1^{i_1} \dots x_n^{i_n}$. For the *lexicographical* ordering, $(i_1, \dots, i_n) \succ (j_1, \dots, j_n)$ (first monomial is *higher* than the second one) if there is $l \leq n$ such that $i_1 = j_1, i_2 = j_2, \dots, i_{l-1} = j_{l-1}$ and $i_l > j_l$. Then the *leading term* $\hat{g}$ of a polynomial g is the monomial which is higher than any other monomial. Now, the reduction is defined as follows. Suppose that the leading term of a given g is divisible by the leading term or some f_i , i.e. $\hat{g} = Q\hat{f}_i$.

²A ring is a set with multiplication and addition. A subset $\mathcal{I}$ of a ring R is called a left (right) ideal if (i) for any $a, b \in \mathcal{I}$ one has $a + b \in \mathcal{I}$ and (ii) for any $a \in \mathcal{I}, c \in R$ one has $ca \in \mathcal{I}$ ($ac \in \mathcal{I}$ respectively). A subset $\{f_i\}$ of $\mathcal{I}$ is a *basis* if any $g \in \mathcal{I}$ equals $\sum r_i f_i$ for some $r_i \in \mathcal{A}$.

Let $g_1 = g - Qf_i$. The leading term of g_1 is lower than the leading term of g and $g_1 \in \mathcal{I}$ if and only if $g \in \mathcal{I}$. One can continue and proceed with g_1 as with g and obtain similarly $g_2, g_3, \dots$. The procedure is repeated until one obtains $g_l \equiv 0$ or an element g_l such that $\hat{g}_l$ is not divisible by any leading term $\hat{f}_i$. One says that g is reduced to g_l modulo the basis $\{f_1, f_2, \dots, f_k\}$.

If a given basis is not a GB one can construct a GB starting from it and using the *Buchberger algorithm*. Suppose that $\hat{f}_i = wq_i$ and $\hat{f}_j = wq_j$ where w, q_i and q_j are monomials and w is not a constant. Define $S(f_i, f_j) = f_i q_j - f_j q_i$. Reduce it modulo $\{f_i\}$ as described above. If the reduction gives a non-zero polynomial add it to the initial basis as f_{k+1} . Consider then such S -polynomials for other pairs of elements (including the new element) and reduce them modulo the ‘current’ basis. If there is nothing to do one obtains a GB. It has been proven by Buchberger [9] that such procedure stops after a finite number of steps.

A classical problem is to find out whether a given element $g \in \mathcal{A}$ is a member of $\mathcal{I}$ or not. This problem can be solved by choosing an ordering and constructing the corresponding GB with the help of the Buchberger algorithm. After that, one applies the reduction procedure modulo the constructed GB to verify whether a given element belongs to the given ideal $\mathcal{I}$. If g is a monomial and $\{g_1, g_2, \dots, g_l\}$ is a GB, then

$$x_1^{a_1} \dots x_n^{a_n} = \sum_{i=1}^l r_i g_i + \sum c_{i_1, \dots, i_n} x_1^{i_1} \dots x_n^{i_n}, \quad (2)$$

where none of the monomials in the second sum is divisible by highest terms of g_i . Since $\{g_1, g_2, \dots, g_l\}$ is a GB, the last sum vanishes if and only if $g \in \mathcal{I}$. If g does not belong to $\mathcal{I}$, the last sum is non-zero, and the part of g belonging to $\mathcal{I}$ is completely included in the first sum.

Let us now denote by $\mathcal{I}$ the (left) ideal generated by the elements f_i which define IBP relations. To solve IBP relations is to express the value of F at an arbitrary point $(a_1, a_2, \dots, a_n)$ in terms of the values of F in a few specially chosen points, i.e. master integrals. This problem can be solved similarly to the algebraic problem described above. Let us think

of the case, where all the indices a_i are positive. Then $F(a_1, a_2, \dots, a_n) = (Y_1^{a_1-1} \dots Y_n^{a_n-1} \cdot F)(1, 1, \dots, 1)$. In this case it is reasonable to consider the operators Y_i as the main operators and get rid of the operators Y_i^{-1} by multiplying (of course, from the left) the operators f_i by sufficiently large powers of the operators Y_i . Let us assume that we are dealing with such f_i .

Let us observe that the situation is quite similar to the above algebraic problem: instead of polynomials in the variables $x_1, \dots, x_n$, we have polynomials in the shift operators $Y_1, \dots, Y_n$. The natural idea is to turn, from the initial basis $\{f_1, \dots, f_i\}$, to a GB $\{g_1, \dots, g_\nu\}$. Indeed, it is known that this can be done similarly to the above case: one introduces an ordering and the notion of the highest term which define the reduction modulo basis, then one can apply a generalization of the Buchberger algorithm. The motivation is the same: this is the GB that characterizes the given ideal in the ‘best’ way, so that the parts which belong to the given ideal do not belong to the second sum in (2). Eventually, one obtains a similar relation, $Y_1^{a_1-1} \dots Y_n^{a_n-1} = \sum_{i=1}^{l'} r_i g_i + \sum c_{i_1, \dots, i_n} Y_1^{i_1-1} \dots Y_n^{i_n-1}$. Let us apply it to F , take the value at $a_i = 1$ and use the fact that the operators of $\mathcal{I}$ give zero on F . We obtain $F(a_1, \dots, a_n) = \sum c_{i_1, \dots, i_n} F(i_1, \dots, i_n)$. Integrals on the right-hand sides of such relations (for various $a_1, \dots, a_n$) are master integrals.

However, any implementation of this similarity, using a generalization of the classical Buchberger algorithm, meets a lot of difficulties. The given problem of solving IBP relations becomes much more complicated at least because one has to consider also non-positive indices a_i .

Another complication is the presence of the variables a_i as non-commutative operators. Moreover, coefficients in monomials in the shift operators, Y_i , can vanish at some points. Still let us imagine a situation where one can apply the Buchberger algorithm to construct a generalization of the GB for solving a reduction problem in the case of positive indices. Simplest examples show that the number of the master integrals associated with this region can be greater than the number of the ‘true’ master integrals.

These complications lead to the natural idea [12] to change the strategy based on GB. For a given family of integrals, $F(a_1, \dots, a_n)$, the whole region for each a_i is decomposed into the regions with $a_i > 0$ and $a_i \leq 0$. The whole region of the multi-indices is decomposed into 2^n sectors $\sigma_\nu = \{(a_1, \dots, a_n) : a_i > 0 \text{ if } i \in \nu, a_i \leq 0 \text{ if } i \notin \nu\}$ labelled by subsets $\nu \subseteq \{1, \dots, n\}$. In a given σ_ν it is natural to consider Y_i for $i \in \nu$ and Y_i^{-1} for other i as basic operators. Another important point in extending Buchberger algorithm, is taking into account boundary conditions, i.e. specify the (*trivial*) sectors where the integrals vanish.

One has to construct [12] a basis of Gröbner type for each non-trivial sector σ_ν . The basic operations are the same as above, i.e. calculating S -polynomials and reducing them modulo current basis, with a chosen ordering. The goal is to construct a so-called *sector* basis [12] (s -basis) which provides the possibility of a reduction to master integrals and integrals of *lower* sectors, i.e. $\sigma_{\nu'}$ for $\nu' \subset \nu$. This point of the strategy is based on multiple examples of solving IBP relations by hand, where one tried to reduce indices to zero. It turns out that, within this strategy, one would construct a true GB only in the case of the sector $\sigma_\emptyset$ but this sector is always trivial.

After constructing s -bases for all non-trivial sectors one obtains a recursive (with respect to the sectors) procedure to evaluate $F(a_1, \dots, a_n)$ at any point and thereby reduce a given integral to master integrals. (See [24] for details of the algorithm.) Examples have shown that the algorithm works at the level of modern calculations. In [12], a non-trivial example of integrals with seven indices was considered and, in [25], reduction problems for two families of HQET integrals with nine indices were solved successfully.

4. Some perspectives

For families of complicated Feynman integrals, a reduction procedure was considered obligatory. On the other hand, it was clear that at least the second strategy of resolving singularities in ϵ within multiple MB representations could be formulated algorithmically and implemented on a computer. (I believe that the first strategy can

also be automated and that the two strategies can be combined in order to achieve an optimization of calculations.) Recently, two algorithmic formulations have appeared [26,27], so that, in complicated situations where one fails to solve the reduction problem by Laporta's algorithm, or by Baikov's method, or by using GB, one can try to calculate every integral, at least numerically, using these formulations.

In fact, the second of these algorithms [27] has been already implemented in *Mathematica*. This code provides a very good precision. For the tennis court integral discussed above this numerical integration provides excellent agreement with the analytic result of [17]. (If such algorithm existed a year ago, the authors of [17] would be satisfied by this powerful check and would not calculate asymptotic behavior when $s/t \rightarrow 0$ by expansion by regions [28] :-)) Anyway, evaluating complicated Feynman integrals, without reduction, becomes now a reliable alternative.

REFERENCES

1. V.A. Smirnov, Phys. Lett. B460 (1999) 397.
2. J.B. Tausk, Phys. Lett. B469 (1999) 225.
3. A.V. Kotikov, Phys. Lett. B254 (1991) 158; B259 (1991) 314; B267 (1991) 123; E. Remiddi, Nuovo Cim. 110A (1997) 1435; T. Gehrmann and E. Remiddi, Nucl. Phys. B580 (2000) 485.
4. K.G. Chetyrkin and F.V. Tkachov, Nucl. Phys. B192 (1981) 159.
5. S. Laporta, Int. J. Mod. Phys. A15 (2000) 5087.
6. T. Gehrmann and E. Remiddi, Nucl. Phys. B601 (2001) 248, 287.
7. C. Anastasiou and A. Lazopoulos, JHEP 0407 (2004) 046.
8. P.A. Baikov, Phys. Lett. B385 (1996) 404; Nucl. Instrum. Methods A389 (1997) 347; V.A. Smirnov and M. Steinhauser, Nucl. Phys. B672 (2003) 199.
9. B. Buchberger and F. Winkler (eds.) *Gröbner Bases and Applications*, (Cambridge University Press, 1998).
10. O.V. Tarasov, Acta Phys. Polon. B29 (1998) 2655; Nucl. Instrum. Meth. A534 (2004) 293.
11. V.P. Gerdt, Nucl. Phys. B (Proc. Suppl.), 135 (2004) 2320; math-ph/0509050; V.P. Gerdt and D. Robertz, cs.SC/0509070.
12. A.V. Smirnov and V.A. Smirnov, JHEP 01 (2005) 001.
13. E.W.N. Glover and M.E. Tejeda-Yeomans, Nucl. Phys. Proc. Suppl. 89 (2000) 196; T. Gehrmann and E. Remiddi, Nucl. Phys. Proc. Suppl. 89 (2000) 251. C. Anastasiou, J.B. Tausk and M.E. Tejeda-Yeomans, Nucl. Phys. Proc. Suppl. 89 (2000) 262.
14. V.A. Smirnov and O.L. Veretin, Nucl. Phys. B566 (2000) 469.
15. C. Anastasiou, T. Gehrmann, C. Oleari, E. Remiddi and J.B. Tausk, Nucl. Phys. B580 (2000) 577;
16. V.A. Smirnov, Phys. Lett. B547 (2002) 239; Phys. Lett. B567 (2003) 193.
17. Z. Bern, L.J. Dixon, and V.A. Smirnov, Phys. Rev. D **72** (2005) 085001.
18. E. Remiddi and J.A.M. Vermaseren, Int. J. Mod. Phys. A15 (2000) 725.
19. V.A. Smirnov, Phys. Lett. B491 (2000) 130; B500 (2001) 330.
20. V.A. Smirnov, Phys. Lett. B524 (2002) 129; V.A. Smirnov, Nucl. Phys. Proc. Suppl. 135 (2004) 252; G. Heinrich and V.A. Smirnov, Phys. Lett. B 598 (2004) 55
21. M. Czakon, J. Gluza and T. Riemann, Acta Phys. Polon. B 36 (2005) 3319; T. Riemann, these proceedings
22. Z. Bern, J.S. Rozowsky, B. Yan, Phys. Lett. B401 (1997) 273; C. Anastasiou, L.J. Dixon, Z. Bern and D.A. Kosower, Phys. Rev. Lett. 91 (2003) 251602.
23. V.A. Smirnov, hep-ph/0209177; *Evaluating Feynman integrals*, (STMP 211; Springer, Berlin, Heidelberg, 2004).
24. A.V. Smirnov, to be published.
25. A.G. Grozin, A.V. Smirnov and V.A. Smirnov, to be published.
26. C. Anastasiou and A. Daleo, hep-ph/0511176.
27. M. Czakon, hep-ph/0511200.
28. M. Beneke and V.A. Smirnov, Nucl. Phys. B522 (1998) 321; V.A. Smirnov, Phys. Lett. B465 (1999) 226; V.A. Smirnov, *Applied Asymptotic Expansions in Momenta and Masses* (Springer, Berlin, Heidelberg, 2001).